

Article title: Pragmatic study of a thromboprophylaxis algorithm in critically ill patients with sars-cov-2 infection

Journal name: Journal of Thrombosis and Thrombolysis

Author names: Maurizio Fattorutto. Yves Bouckaert. Jonathan Brauner. Stéphane Franck. Fabrice Bouton. Danielle Heuse. Charlotte Bouckaert. Arnaud Bruyneel.

Corresponding author: Maurizio Fattorutto, Department of Anesthesiology, Centre Hospitalier Universitaire Tivoli, Avenue Max Buset 34, 7100 La Louvière, Belgium. E-mail address: maurizio.fattorutto@chu-tivoli.be

Table 5. ROTEM™ viscoelastic analyses and thrombotic/inflammatory markers

MCF-EXTEM™	Fibrinogen	D-dimer	WBC	CRP
(mm)	(g/L)	(ng/mL FEU)	(x10³ cells/μL)	(mg/L)
80	6	5000	4.2	89.6
79	7.2	816	8.4	80.8
82	7.3	876	8.3	90
77	7	912	6.8	75.1
75	4.5	1558	14.8	9.9
69	4.3	415	8.5	151.5
77	7.4	699	12.4	225.1
80	4.4	2027	12.4	21.5
75	6.6	904	10.3	212

Each of the analyses presented in the table was performed on the same blood sample at a time when each of the patients presented with severe ARDS but none was in septic shock. None of these samples was taken near (+/- 24 hours) a thromboembolic event or a fatal outcome. Abbreviations: CRP: C-reactive protein; FEU: fibrinogen equivalent units; MCF: Maximum Clot Firmness (symbolize the final viscoelastic property of the thrombus); WBC: white blood cell count

